

LiDCO Science & Studies Bibliography

L1.1 Linton RAF, Band DM & Haire KM. (1993) A new method of measuring cardiac output in man using lithium dilution. *British Journal of Anaesthesia*; 71: 262-266

L1.2. Band D, Linton R. (1994) A New way of Measuring Cardiac Output. *The Physiological Society Magazine* 17:21

L1.3. Linton R, Band D, O'Brien T, Jonas MM & Leach R. (1997) Lithium dilution cardiac output measurement: A comparison with thermodilution. *Critical Care Medicine*; 25: 1796-1800

L1.4. Kurita T, Morita K, Kato S, Kikura M, Horie M, Ikeda K. (1997) Comparison of the accuracy of the lithium dilution technique with the thermodilution technique for measurement of cardiac output. *British Journal of Anaesthesia*; 79: 770-775

L1.5. Linton RAF, Band DM, O'Brien TK, Linton NWF & Jonas MM. (1998) Lithium dilution cardiac output measurement - a brief review. In: *State-of-the-Art Technology in Anesthesia and Intensive Care* , pp 61-66. Edited by: Ikeda K, Doi M & Kazama T; pub. Elsevier Science B.V., Amsterdam

L1.6. Jonas MM, Kelly FE Linton RAF, Band DM, O'Brien TK, Linton NWF. (1999) A comparison of lithium dilution cardiac output measurements made using central and antecubital venous injection of lithium chloride. *Journal of Clinical Monitoring & Computing*, 15: 525-528

L1.7. Jonas, Linton, O'Brien, Band, Linton, Kelly, Burden, Chevalier, Thompson, Birch and Powell. (2001) The pharmacokinetics of intravenous lithium chloride in patients and normal volunteers. *Journal of Trace and Microprobe Techniques*. 19: 313-320

L1.8. Garcia-Rodriguez C, Pittman J, Cassell CH, Sum-Ping J, El-Moalem H, Young C, Mark JB. (2002) Lithium dilution cardiac output measurement: A Clinical assessment of central venous and peripheral venous indicator injection. *Crit Care Med* Vol. 30, No. 10 p2199-2204

L1.9. Kurita T, Morita K, Kawasaki H, Fujii K, Kazama T, Sato S. (2002) Lithium Dilution Cardiac Output Measurement in Oleic Acid-Induced Pulmonary Edema. *Journal of Cardiothoracic and Vascular Anesthesia*, Vol 16, No 3 (June): pp 334-337

L1.10. Mason D, O'Grady M, Woods P, McDonnell W. (2002) Effect of background serum lithium concentrations on the accuracy of lithium dilution cardiac output determination in dogs. *American Journal of Veterinary Research*; 63 : 1048 – 1052

L1.11. Pittman J, Bar Yosef S, SumPing J, Sherwood M, Mark J (2005) Continuous cardiac output monitoring with pulse contour analysis: A comparison with lithium indicator dilution cardiac output measurement. *Crit Care Med*. 33 (9) 2015-2021

L1.12. Cecconi M, Dawson D, Grounds R, Rhodes A. (2008) Lithium dilution cardiac output measurement in the critically ill patient: determination of precision of the technique. *Intensive Care Medicine*; 35 : 498 – 504

L1.13. Kemps H, Thijssen E, Schep G, Sleutjes B, De Vries W, Hoogeveen A, Wijn P, Doevendans P. (2008) Evaluation of two methods for continuous cardiac output assessment during exercise in chronic heart failure patients. *J Appl Physiol* 105: 1822-1829

L1.14. Hannegard-Hamrin T, Odeberg-Wenerman S (2010) Lithium indicator technique for cardiac output in children: reproducibility of the calibration factor. Proceedings of the 2010 Annual Meeting of the ESICM

Further Presented & Published LiDCO System Studies

L2.1. Linton RAF, Linton NWF & Band DM. (1995) A new method of analysing indicator dilution curves. *Cardiovascular Research*; 30: 930-938

L2.2. Morita K, Kurita T, Kikura M, Ikeda K. (1995) Accuracy of the Lithium Dilution Cardiac Output Measurement. Meeting of the European Society for Computing and Technology in Anaesthesia and Intensive Care, Palermo

L2.3. Linton R, Band D. (1995) Cardiac Output Measured by Indicator Dilution Using Lithium Chloride. Meeting of the Japanese Society for Anaesthesia, Hamamatsu

L2.4. Band DM, Linton RAF, O'Brien TK, Jonas M & Linton NWF. (1997) The shape of indicator dilution curves used for cardiac output measurement in man. *Journal of Physiology*; 498: 225-229

L2.5. Linton R, Turtle M, Band D, O'Brien T & Jonas M. (1998) In vitro evaluation of a new lithium dilution method of measuring cardiac output and shunt fraction in patients undergoing venovenous extracorporeal membrane oxygenation. *Critical Care Medicine*; 26: 174-177

L2.6. Linton RAF, Taylor PM, Linton NWF, Flach EJ, O'Brien TK, Band DM. (1999) Cardiac output measurement in an anaesthetised giraffe. *The Veterinary Record*, October 23;145(17):498-9

L2.7. Mappes A, Wiersich S, Siniawski H, Lindert J, Kuppe H. (1999) Comparison of three different technologies of measuring cardiac output (CO) in patients undergoing open heart surgery. Meeting of the American Society of Anesthesiology, Dallas October

L2.8. Peruzzi W. (2000) Ion selective electrodes and critical care monitoring. *American Society of Critical Care Anesthesiologists Interchange*. 12: 3&7

L2.9. Mappes A, Wiersich S, Koster A, Lindert J, Kuppe H. (1999) Lithium dilution cardiac output measurement in patients undergoing coronary artery bypass grafting. Meeting of the European Society of Intensive Care Medicine, Berlin, October

L2.10. Linton RAF, Turtle M, Band DM, O'Brien TK, Jonas MM & Linton NWF. (1999) A new technique for measuring cardiac output and shunt fraction during venovenous extracorporeal membrane oxygenation. *Perfusion*; 14: 43-47

L2.11. Garcia-Rodriguez C, Cassell C, Young C, Sum Ping J. Mark J. (2000) Cardiac output measurement without pulmonary artery or central venous catheterization: A clinical assessment of the lithium dilution method. *American Society of*

Anesthesiology, San Francisco October

L2.12. Linton RAF, Linton NWF, Kelly F. (2000) How accurate is clinical assessment of cardiac output in the early postoperative period following cardiac surgery? American Society of Anesthesiology, San Francisco CA October

L2.13. Linton RA, Jonas MM, Tibby SM, Murdoch IA, O'Brien TK, Linton NW, Band DM. (2000) Cardiac output measured by lithium dilution and transpulmonary thermodilution in patients in a paediatric intensive care unit. Intensive Care Med. Oct; 26(10): 1507-11. [*The use of the LiDCO lithium dilution measurement of cardiac output is not licensed for use in patients of less than 40 kgs / 88lbs in weight*]

L2.14. Linton RA, Young LE, Marlin DJ, Blissett KJ, Brearley JC, Jonas MM, O'Brien TK, Linton NW, Band DM Jones RS. (2000) Cardiac output measured by lithium dilution, thermodilution and transesophageal Doppler echocardiography in anesthetized horses. American Journal of Veterinary Research Jul; 61(7): 731-7

L2.15. Linton RAF. (2000) Cardiac output measured by lithium dilution. EACTA meeting, Aarhus Denmark June

L2.16. Mappes A, Gruendel M, Lindert J, Wiersich S, Kuppe H. (2000) Hemodynamic monitoring in cardiac surgery: changes in mean arterial pressure do not correlate with the associated changes in cardiac output. EACTA meeting, Aarhus Denmark June

L2.17. Mappes A, Lindert J, Wiersich S, Gruendel M, Kuppe, H. (2000) Hemodynamic monitoring during cardiac surgery: the new LiDCO method. Cardiothoracic Technologies and Techniques meeting, Bal Harbour Florida, January

L2.18. Mason DJ, O'Grady M, Woods JP, McDonell W. (2001) Assessment of lithium dilution cardiac output as a technique for measurement of cardiac output in dogs. American Journal of Veterinary Research Aug; 62(8): 1255-1261

L2.19. Hatfield C, McDonell W, Lemke D, Black W. (2001) Pharmacokinetics and toxic effects of lithium chloride after intravenous administration in conscious horses. American Journal of Veterinary Research Sep; 62(9): 1387-1392

L2.20. Siddall VJ, Gould R, Peruzzi W. (2001) Comparison of clinician assessment of haemodynamic status with actual data obtained through a minimally invasive lithium dilution method. Presented at the 47th Annual International Respiratory Congress San Antonio, Texas

L2.21. Jonas M, Bruce R, Knight J, Kelly F, O'Brien T, Band D. (2001) Comparison of cardiac output measurements using a continuous arterial waveform analysis monitor (PulseCO) with an indicator dilution technique (LiDCO) in patients in intensive care. Presented at the ICCM Sydney, October

L2.22. Mason DJ, O'Grady M, Woods JP, McDonell W (2002) Comparison of a central and a peripheral (cephalic vein) injection site for the measurement of cardiac output using the lithium-dilution cardiac output technique in anesthetized dogs. Can J Vet Res. 66(3): 207-210

L2.23. Corley K, Donaldson L, Furr M. (2002) Comparison of lithium dilution and thermodilution cardiac output measurements in anaesthetised neonatal foals. *Equine Vet J* 2002 Sept;34 (6) 598-601

L2.24. Jonas M, Tanser S. (2002) Lithium dilution measurement of cardiac output and arterial pulse waveform analysis: an indicator dilution calibrated beat-by-beat system for continuous estimation of cardiac output. *Current Opinion in Critical Care*; 8 : 257 – 261

L2.25. P.R. Roberts, S. Allen, S. Robinson, S.J. Tanser, M.M. Jonas, J.M. Morgan. (2002) Use of lithium dilution assessment of cardiac output to optimise right/left ventricular activation In Resynchronisation Therapy Presented at: 80th Anniversary British Cardiac Society Annual Scientific Conference Harrogate. Pub. In *Heart*, May 2002: 87 Supplement II (Abstract 146)

L2.26. Mappes A, Dittrich S, Gruendel M, Huebler M, Hetzer R, Kuppe H. Lithium-based indicator dilution measurement of cardiac output in a 10 year old child after orthotopic heart transplantation. Case Report- Berlin Heart Center. [*The use of the LiDCOplus Hemodynamic Monitor is unlicensed in patients of less than 40 kgs / 88lbs in weight.*]

L2.27. Corley KTT, Donaldson LL, Durando MM, Birks EK. (2003) Cardiac output technologies with special reference to the horse. *Journal Vet Intern Med*. May-Jun;17(3):262-72

L2.28. Hallowell G, Corley K (2005) Use of lithium dilution and pulse contour analysis cardiac output determination in anestheized horses: a clinical evaluation. *Vet Anaes & Analgesia* 32, 201-211

L2.29. Lobo S, Oliveira N, Lobo F, Rezende E, Borges B, Cunrath G, Silva J, Ronchi L (2007) Oxygen delivery optimization using lithium indicator dilution and pulse power analysis during major surgery in high-risk patients. Poster Presentation ISICEM, Brussels

L2.30. Durando M, Corley K, Boston R, Birks E. (2008) Cardiac output determination by use of lithium dilution during exercise in horses. *American Journal Veterinary Research*; 69 : 1054 – 1060

L2.31. Flood S, Bodenham A (2010) Lithium: mimicry, mania and muscle relaxants Continuing Education in Anaesthesia, Critical Care & Pain April 15 2010 doi10.1093/bjaceaccp/mkq008

L2.32 Valverde A, Gianotti G, Rioja E, Hathway A. (2011) Comparison of cardiac output determined by arterial pulse pressure waveform analysis method (Flotrac/Vigileo) versus lithium dilution method in anesthetized dogs. *J Veterinary Emergency and Critical Care* vol 21:328-334

L2.33 Shih A, Maisenbacher H, Bandt C, Ricco C, Bailey J, Rivera J, Estrada A. (2011) Assessment of cardiac output measurement in dogs by transpulmonary pulse contour analysis. *Journal of Veterinary Emergency and Critical Care* vol 21

(4):321-327

L2.34 Shih A, Giguere S, Vigani A, Shih R, Thuramalla N, Bandt C. (2011) Determination of cardiac output velocity dilution in normovolemia and hypovolemia in dogs. *Veterinary Anaesthesia and Analgesia* vol 38(4): 279-285

P1.1 Hamilton TT, Huber LM, Jessen ME. (2002) PulseCO: A Less-Invasive Method to Monitor Cardiac Output From Arterial Pressure After Cardiac Surgery. *Ann Thorac Surg*; 74:S1408-12

P1.3 Pearse R, Dawson D, Fawcett J, Rhodes A, Grounds RM, Bennett ED (2005) Early goal-directed therapy after major surgery reduces complications and duration of hospital stay. A randomised, controlled trial. *Crit Care* 9 (6) 687-693

P1.4 Pittman J, Bar Yosef S, SumPing J, Sherwood M, Mark J (2005) Continuous cardiac output monitoring with pulse contour analysis: A comparison with lithium indicator dilution cardiac output measurement. *Crit Care Med*. 33 (9) 2015-2021

P1.5 Belloni L, Pisano A, Natale A, Piccirillo M, Piazza L, Ismeno G, De Martino G (2007) Assessment of Fluid-Responsiveness Parameters for Off-Pump Coronary Artery Bypass Surgery: A Comparison Among LiDCO, Transesophageal Echocardiography, and Pulmonary Artery Catheter. *J Cardiothorac Vasc Anesth*. 2008 Apr;22(2):243-8.

P1.7 Costa MG, Della Rocca G, Chiarandini P, Mattelig S, Pompei L, Barriga MS, Reynolds T, Cecconi M, Pietropaoli P (2007) Continuous and intermittent cardiac output measurements in hyperdynamic conditions: pulmonary artery catheter versus lithium dilution technique. *Intensive Care Medicine* DOI 10.1007/s00134-007-0878-6

P1.8 Missant C, Rex S, Wouters P. (2008) Accuracy of cardiac output measurements with pulse contour analysis (PulseCO) and Doppler echocardiography during off-pump coronary artery bypass grafting. *European Journal of Anaesthesiology* 25(3): 243-248

P1.9 Kemps H, Thijssen E, Schep G, Sleutjes B, De Vries W, Hoogeveen A, Wijn P, Doevendans P. (2008) Evaluation of two methods for continuous cardiac output assessment during exercise in chronic heart failure patients. *Journal of Applied Physiology* 105: 1822-1829

P1.10 Marquez J, McCurry K, Severyn D, Pinsky M. (2008). Ability of Pulse Power, Esophageal Doppler and Arterial Pulse Pressure to Estimate Rapid Changes in Stroke Volume in Humans. *Crit Care Med* 36(11) 3001 – 3007

P1.11 Mills E, Jonas M, Wolff C, O'Brien T (2010) PulseCO consistency: variation in calibration factor over 24 and 48 hours. Poster presentation ISICEM, Brussels. *Critical Care* 14(1), p101

P1.12 Jonas M, Mills E, Wolff C, O'Brien T (2010) Effect of cardiac arrhythmias on PulseCO calibration and performance. Poster presentation ISICEM, Brussels. *Critical Care* 14(1), p102

P1.13 Green D, Paklet L (2010) Latest developments in peri-operative monitoring of

the high-risk major surgery patient. *International Journal of Surgery* 8 90-99

P1.14 Abdel-Galil K, Craske D, McCaul J (2010) Optimisation of intraoperative haemodynamics: early experience of its use in major head and neck surgery. *Brit Journal of Oral and Maxillofacial Surgery* 48 (3) 189-191

P1.17 Purushothaman B, O'Brian T, Green D (2010) The Hemodynamic Effects of Anaesthetic Induction and Their Correlation with Changes in Depth of Anaesthesia. *Proceedings of the 2010 Annual Meeting of the ASA*

P1.18 Koff M, Richard K, Novak M, Canneson M, Dodds T (2010) Elevated PPV Predict an increased Length of Stay and Morbidity during High Risk Abdominal Surgery. *Proceedings of the 2010 Annual Meeting of the ASA*

P1.19 Hadian M, Severyn D, Pinsky M (2010) The effects of vasoactive drugs on pulse pressure and stroke volume variation in postoperative ventilated patients. *Critical Care In press*

P2.11 Linton N, Linton R (2003) Haemodynamic response to a small intravenous bolus injection of epinephrine in cardiac surgical patients. *European Journal of Anaesthesiology* 20: 298-304

P2.14 Kirwan C, Smith J, Lei K, Beale R (2005) A comparison of two calibrated continuous arterial pressure waveform based measurements of cardiac output over 24 hour. *Crit Care Med.* 33(12) Suppl; 208-S; A56

P2.18 De Wilde RBP, Schreuder JJ, van den Berg PCM, Jansen JRC (2007) An evaluation of cardiac output by five arterial pulse contour techniques during cardiac surgery. *Anaesthesia*: 62, 760-768

P2.22 Dyer RA, Piercy JL, Reed AR, Lombard CJ, Schoeman LK, James MF (2008) Hemodynamic changes associated with spinal anesthesia for cesarean delivery in preeclampsia. *Anesthesiology* Vol. 108, No. 5, 802 – 811

P2.24 Jonas M, Fennell J, Brudney CS. (2008). Haemodynamic optimisation of the surgical patient revisited. *Anaesthesia International*, Spring 2008 Vol 2 No1

P2.25 Archer T, Knape K, Liles D, Wheeler A, Carter B. (2008). The hemodynamics of oxytocin and other vasoactive agents during neuraxial anesthesia for cesarean delivery: findings in six cases. *Int. J. Obstetric Anesthesia* Vol. 17:3, 247-254.

P2.27 Kim H, Hadian M, Severyn D, Pinsky M (2009). Cross-comparison of the trending accuracy of continuous cardiac output measurement devices in postoperation patients. Poster presentation ISICEM, Brussels. *Critical Care*, 13(Suppl 1):P209doi:10.1186/cc7373

P2.47 Dyer R, Reed A, van Dyk D, Arcache M, Hodges O, Lombard C, Greenwood J, James M (2009) Hemodynamic Effects of Ephedrine, Phenylephrine, and the Coadministration of Phenylephrine with Oxytocin during Spinal Anesthesia for Elective Cesarean Delivery. *Anesthesiology* 111:753-65

P2.48 Cecconi M, Monti G, Dawson D, Vamadan S, Hamilton M, Della Rocca G,

Grounds RM, Rhodes A (2009) Dynamic Indices by PulseCO in post surgical intensive care patients. Proceedings of the 2009 Annual Meeting of the ESICM

P2.62 P Brass, E Mills, J Latza, J Peters, E Berendes. LiDCOrapid and PICCOplus preload response parameter validation study. 2011 doi:10.1186/cc9481

P2.63 P Brass, E Mills, J Latza, J Peters, E Berendes. Comparison of cardiac index: LiDCOrapid and PICCOplus in the ICU. 2011 doi:10.1186/cc9482

P2.65 C Willars, A Dada, D Green, Functional haemodynamic monitoring the relative merits of SVV, SVP and PPV as measured by the LiDCOrapid in predicting fluid responsiveness in high-risk surgical patients. Poster presentation, 2011 doi:10.1186/cc9486

P2.67. Langesaeter E, Rosseland L, Stubhaug A (2011). Haemodynamic effects of oxytocin in women with severe preeclampsia. *Intl Jnl of Obstetric Anaesthesia* 20 (1) 26 – 29

P2.68. Langesaeter E, Dyer R (2011). Maternal haemodynamic changes during spinal anaesthesia for caesarean section. *Current Opinion in Anaesthesiology* 24 (3) 242 - 248

P2.69. Passos R, Batista P (2011) Fluid removal in critically ill patients during hemodialysis: is there a role for functional hemodynamic monitoring? Proceedings of the 6th International Symposium on Intensive Care and Emergency Medicine for Latin America 15 (suppl 2): P30 doi:10.1186/cc10178

P2.70 Eide P (2011) Cardiac output in idiopathic normal pressure hydrocephalus: association with arterial blood pressure and intracranial pressure wave amplitudes and outcome of shunt surgery. *Fluids and Barriers of the CNS* 8 (11) 1-10

P2.71 Geerts B, Aarts L, Groeneveld A, Jansen J. Predicting cardiac output responses to passive leg raising by a PEEP-induced increase in central venous pressure, in cardiac surgery patients. *British Journal of Anaesthesia* 107 (2): 150-6

P2.72 Singh S, Kuschner W, Lighthall G. (2011) Perioperative intravascular fluid assessment and monitoring: a narrative review of established and emerging technologies. *Anesthesia Research and Practice* Article ID 231493, doi:10.1155/2011/231493

P2.73 Hamilton M, Cecconi M, Rhodes A. (2011) A systematic review and meta-analysis on the use of pre-emptive hemodynamic intervention to improve postoperative outcomes in moderate and high risk surgical patients. *Anesthesia and Analgesia* June vol. 112:(6) 1392 -1402

P2.74 Lee A, Cohn J, Ranasinghe S. (2011) Cardiac output assessed by invasive and minimally invasive techniques. *Anesthesia Research and Practice* Article ID 475151, doi:10.1155/2011/475151

P2.75 Geerts B, Aarts L, Jansen J. (2011) Methods in pharmacology: measurement

of cardiac output. British Journal of Clinical Pharmacology vol 71 (3): 316 – 330

P2.76 Potocnik I, Jankovic V, Kremzar B, Stupnik T. (2011) Haemodynamic change after induction of anaesthesia with sevoflurane vs. propofol. Signa Vitae vol 6(2): 52-57

P2. 77 Mora B, Ince I, Birkenbreng B, Skhirtladze K, Pernicka E, Ankersmit H, Dworschak M. (2011) Validation of cardiac output measurement with the LiDCO pulse contour system in patients with impaired left ventricular function after cardiac surgery. Anaesthesia vol 66(8): 675-681

P2.78 De Santis P, Marano C, Cavallaro F, Dell'Anna A, Santis P, Bonarrigo C, Falcone C, Sandroni C. (2011) Crit Care 15(Suppl 1):p64

P2.79 Alhashemi A, Cecconi M, Hofer C. (2011) Cardiac output monitoring: an integrative perspective. Critical Care 15:214

P2.80 Chalmers A, Turner M, Anand R, Puxeddu R, Brennan P. (2011) Cardiac output monitoring to guide fluid replacement in head and neck microvascular free flap surgery – what is current practice in the UK? British Journal of Oral and Maxillofacial Surgery doi:10.1016/j.bjoms.2011.08.010

LiDCO.com website clinical papers (May 2017)

[Cannesson M, Gan TJ. PRO: Perioperative Goal-Directed Fluid Therapy is an essential element of an Enhanced Recovery Protocol. Anesth Analg. 2016;122\(5\):1258-60.](#)

[Mosquera C, Koutlas NJ, Fitzgerald TL. A single surgeon's experience with Enhanced Recovery after Surgery: An Army of One. Am Surg. 2016;82\(7\):594-601.](#)

[Tengberg LT, Bay-Nielsen M, Bisgaard T, Cihoric M, Lauritsen MI, Foss NB. Multidisciplinary perioperative protocol in patients undergoing acute high-risk abdominal surgery. Br J Surg. 2017 Mar;104\(4\):463-471.](#)

[Hata J, Stotts C, Shelsky C, Bayman E, Frazier A, Wang J, Nickel E. Reduced mortality with noninvasive hemodynamic monitoring of shock. J Crit Care. 2011;26\(2\):224.E1-8.](#)

[Pearse R, Dawson D, Fawcett J, Rhodes A, Grounds RM, Bennett ED. Early goal-directed therapy after major surgery reduces complications and duration of hospital stay. A randomised, controlled trial. Crit Care. 2005;9 \(6\):687-693 & Ebm C, Cecconi M, Sutton L, Rhodes A. A Cost-Effectiveness Analysis of Postoperative Goal-Directed Therapy for High-Risk Surgical Patients. Crit Care Med. 2014. DOI: 10.1097/CCM000000000000164](#)

[Jain A & Dutta A Stroke Volume Variation as a Guide to Fluid Administration in Morbidly Obese Patients Undergoing Laparoscopic Bariatric Surgery. Obes Surg. 2010. DOI 10.1007/s11695-009-0070](#)

[Wei Xiao, Qing-Fang Duan, Wen-Ya Fu,1 Xin-Zuo Chi, Feng-Ying Wang, Da-Qing Ma, Tian-Long Wang, and Lei Zhao. Goal-directed Fluid Therapy May Improve](#)

[Hemodynamic Stability of Parturient with Hypertensive Disorders of Pregnancy Under Combined Spinal Epidural Anesthesia for Cesarean Delivery and the Well-being of Newborns. Chin Med J \(Engl\). 2015 ;128\(14\):1922–1931. doi: 10.4103/0366-6999.160546](#)

[Bisgaard J, Gilsaa T, Ronholm E, Toft P. Haemodynamic optimisation in lower limb arterial surgery: Room for improvement? Acta Anaesthesiol Scand. 2013;57:189-198.](#)

[Green D, Bidd H, Rashid H. Multimodal intraoperative monitoring: An observational case series in high risk patients undergoing major peripheral vascular surgery. Int J Surg \(2014\);12\(3\):231-6. doi: 10.1016/j.ijvsu.2013.12.016. Epub 2014 Jan 8.](#)

[Han G, Liu K, Xue H, Zhao P. Application of LiDCO-Rapid in peri-operative fluid therapy for aged patients undergoing total hip replacement. Int J Clin Exp Med 2016;9\(2\):4473-4478, www.ijcem.com /ISSN:1940-5901/IJCEM0010819](#)

[Lobo S, et al. Restrictive strategy of intraoperative fluid maintenance during optimization of oxygen delivery decreases major complications after high-risk surgery. Crit Care. 2011;15:R226. doi:10.1186/cc10466](#)

[Osawa EA, Rhodes A, Landoni G, Galas FR, et al. Effect of Perioperative Goal-Directed Hemodynamic Resuscitation Therapy on Outcomes Following Cardiac Surgery: A Randomized Clinical Trial and Systematic Review. Crit Care Med. 2016;44\(4\):724-33. doi: 10.1097/CCM.0000000000001479.](#)

[Fitzgerald T, Mosquera C, Koutlas N, Vohra N, et al. Enhanced recovery after surgery in a single high-volume surgical oncology unit: Details matter. Presented at the 11th Annual Academic Surgical Congress \(ASC\), Jacksonville, Florida, Feb 2016](#)

[Karran A, Wheat J, Chan D, Blake P, et al. Propensity Score Analysis of an Enhanced Recovery Programme in Upper Gastrointestinal Cancer Surgery. World J Surg. 2016;40:1645–1654. DOI 10.1007/s00268-016-3473-6A.](#)

[Preston SR, Markar SR, Baker CR, Soon Y, et al. Impact of multidisciplinary standardized clinical pathway on perioperative outcomes in patients with oesophageal cancer. Br J Surg. 2013;100\(1\):105-12. DOI: 10.1002/bjs.8974](#)

[Pearse RM, Harrison DA, MacDonald N, Michael A, Gillies MA, Blunt M, Ackland G, Grocott MP, Ahern A, Griggs K, Scott R, Hinds C, Rowan K for the OPTIMISE study group. Effect of a peri-operative, cardiac output-guided, hemodynamic therapy algorithm on outcomes following major gastrointestinal surgery: A randomized clinical trial and updated systematic review. JAMA. 2014;311\(21\):2181-2190. doi:10.1001/jama.2014.5305.](#)

[Miller T, Thacker J, White W, Mantyh C, et al. Reduced length of hospital stay in colorectal surgery after implementation of an enhanced recovery protocol. Anesth Analg 2014;118:1052–61.](#)

[Dunne DF, Yip VS, Jones RP, McChesney EA, et al. Enhanced recovery in the resection of colorectal liver metastases. J Surg Oncol. 2014. doi:10.1002/jso.23616](#)

Jones C, Kelliher L, Dickinson M, Riga A, Worthington T, Scott M J, Vandrevala T, Fry C H, Karanjia N and Quiney N. Randomized clinical trial on enhanced recovery versus standard care following open liver resection. Br J Surg 2013 Jul;100(8):1015-24. Doi: 10.1002/bjs.9165

