# How to save lives in Sepsis

LiDCO hemodynamic monitoring reduces mortality in Sepsis



**GLOBAL PRIORITY** 

respond to fluids





"It is likely that morbidity and Portality in 5ex








**Management** 





**Calibrate** 

**Minimally Invasive** 





# Clinical papers showing improved outcome

# Sepsis | Reduced mortality with noninvasive hemodynamic monitoring of shock

# **Patient Population**

ICU shock patients

### **LiDCO Monitor**

LiDCOplus

# Trial Design

Observational study comparing no hemodynamic monitoring vs pulmonary artery vs LiDCOplus managed shock patients

# **Outcome Impact**

Treatment of patients using the LiDCOplus monitor significantly reduced the observed mortality rate to 13% against 32% and 20% in the invasively monitored and 37% in the unmonitored patient groups

<sup>5</sup> Hata J, Stotts C, Shelsky C, Bayman E, Frazier A, Wang J, Nickel E. Reduced mortality with noninvasive hemodynamic monitoring of shock. J Crit Care. 2011;26(2):224.E1-8.

# A rational approach to fluid therapy in sepsis

## **Patient Population**

Severe sepsis and septic shock

## **Purpose**

The purpose of this consensus is to provide support to the bedside clinician regarding the diagnosis, management and monitoring of shock

# L

#### Methods

The European Society of Intensive Care Medicine (ESICM) formed a task force of 12 experts

#### Results

44 consensus statements that can be used at the bedside to diagnose, treat and monitor patients with shock

- 23 Marik P, Bellomo R. A rational approach to fluid therapy in sepsis. Br J Anaesth. 2016;116(3):339-49.
- <sup>1</sup> Reinhart, K., Daniels, R., Kissoon, N. et al. Recognizing Sepsis as a Global Health Priority A WHO Resolution. N Engl J Med 2017; 377:414-41
- Cecconi, M., De Backer, D., Antonelli, M. et al. Intensive Care Med (2014) 40: 1795. doi:10.1007/s00134-014-3525-z



Contact us on how LiDCO fits into your Sepsis pathway



